

# DR ACADEMY

## DO RIGHT FOR GENUINE EDUCATION

KCET EXAMINATION – 2020

SUBJECT : CHEMISTRY

DATE :- 31-07-2020

TIME : 02.30 PM TO 03.50 PM

1. Copper is extracted from copper pyrites by  
 a) Thermal decomposition  
 b) Reduction by coke  
 c) Electrometallurgy  
 d) Auto reduction

Ans. d

2. Function of potassium ethyl xanthate in froth floatation process is to make the ore  
 a) Lighter  
 b) Hydrophobic  
 c) Hydrophilic  
 d) Heavier

Ans. b

3. Sulphide ore on roasting gives a gas X. X reacts with  $\text{Cl}_2$  in the presence of activated charcoal to give Y. Y is:  
 a)  $\text{SO}_2\text{Cl}_2$     b)  $\text{S}_2\text{Cl}_2$     c)  $\text{SCl}_6$     d)  $\text{SOCl}_2$

Ans. a

4. Aqueous solution of a salt (A) forms a dense white precipitate with  $\text{BaCl}_2$  solution. The precipitate dissolves in dilute  $\text{HCl}$  to produce a gas (B) which decolourises acidified  $\text{KMnO}_4$  solution

A and B respectively are:

- a)  $\text{BaSO}_3, \text{SO}_2$     b)  $\text{BaSO}_4, \text{H}_2\text{S}$   
 c)  $\text{BaSO}_3, \text{H}_2\text{S}$     d)  $\text{BaSO}_4, \text{SO}_2$

Ans. a

5. Bond angle in  $\text{PH}_4^+$  is more than that of  $\text{PH}_3$ . This is because  
 a) Lone pair – bond pair repulsion exists in  $\text{PH}_3$   
 b)  $\text{PH}_4^+$  has square planar structure  
 c)  $\text{PH}_3$  has planar trigonal structure  
 d) Hybridisation of P changes when  $\text{PH}_3$  is converted to  $\text{PH}_4^+$

Ans. a

6. Incorrectly matched pair is:  
 a)  $\text{XeO}_3$  – pyramidal  
 b)  $\text{XeF}_4$  – tetrahedral  
 c)  $\text{XeF}_6$  – distorted octahedral  
 d)  $\text{XeOF}_4$  – square pyramidal

Ans. b

7. Phosphorus pentachloride  
 a) On hydrolysis gives an oxo acid of phosphorus which is tribasic  
 b) On hydrolysis gives an oxo acid of phosphorus which is a good reducing agent  
 c) Has all the five equivalent bonds  
 d) Exists as an ionic solid in which cation has octahedral structure and anion has tetrahedral structure

Ans. a

8. Identify the set of paramagnetic ions among the following:  
 a)  $\text{V}^{2+}, \text{Co}^{2+}, \text{Ti}^{4+}$     b)  $\text{Ni}^{2+}, \text{Cu}^{2+}, \text{Zn}^{2+}$   
 c)  $\text{Ti}^{3+}, \text{Cu}^{2+}, \text{Mn}^{3+}$     d)  $\text{Sc}^{3+}, \text{Ti}^{3+}, \text{V}^{3+}$

Ans. c

9. How many moles of acidified  $\text{K}_2\text{Cr}_2\text{O}_7$  is required to liberate 6 moles of  $\text{I}_2$  from an aqueous solution of  $\text{I}^-$ ?  
 a) 2    b) 1    c) 0.25    d) 0.5

Ans. a

10.  $\text{Cu}_2\text{Cl}_2$  and  $\text{CuCl}_2$  in aqueous medium  
 a)  $\text{CuCl}_2$  is more stable than  $\text{Cu}_2\text{Cl}_2$   
 b) Stability of  $\text{Cu}_2\text{Cl}_2$  is equal to stability of  $\text{CuCl}_2$   
 c) Both are unstable  
 d)  $\text{Cu}_2\text{Cl}_2$  is more stable than  $\text{CuCl}_2$

Ans. a

11. The Co-ordination number of Fe and Co in the complex ions,  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$  and  $[\text{Co}(\text{SCN})_4]^{2-}$  are respectively:  
 a) 3 and 4    b) 6 and 8  
 c) 4 and 6    d) 6 and 4

Ans. d

12. Number of stereoisomers exhibited by  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$  is  
 a) 4    b) 2    c) 5    d) 3


Ans. d

13. Give the IUPAC name of  $[\text{Pt}(\text{NH}_3)_4][\text{PtCl}_4]$  is
- Tetra ammine platinum (o) tetra chlorido platinum (IV)
  - Tetra ammine palatinate (II) tetra chlorido platinum (II)
  - Tetra ammine palatinate (o) tetra chlorido platinum (IV)
  - Tetra ammine platinum (II) tetra chlorido palatinate (II)

**Ans. d**

14. Prolonged exposure of chloroform in humans may cause damage to liver. It is due to the formation of the following compound
- $\text{CCl}_4$
  - $\text{COCl}_2$
  - $\text{CH}_2\text{Cl}_2$
  - $\text{Cl}_2$

**Ans. b**



15. Which of the following halide shows highest reactivity towards  $\text{S}_{\text{N}}1$  reaction?
- $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$
  - $\text{CH}_3 - \text{CH}_2\text{Cl}$
  - $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2\text{I}$
  - $\text{C}_6\text{H}_5\text{Cl}$

**Ans. a**

**Sol.** Rate of  $\text{S}_{\text{N}}1$  reaction is directly proportional to stability of carbocation or Reactivity of  $\text{S}_{\text{N}}1$  reaction is influenced by stability of carbocation.


16. In the reaction



The number of possible isomers for the organic compound X is

- 4
- 5
- 3
- 2

**Ans. d**



x = isobutane and it has two isomers.

17. Which of the following on heating gives an ether as major products?
- P:  $\text{C}_6\text{H}_5\text{CH}_2\text{Br} + \text{CH}_3\text{ONa}$   
 Q:  $\text{C}_6\text{H}_5\text{ONa} + \text{CH}_3\text{Br}$   
 R:  $(\text{CH}_3)_3\text{C} - \text{Cl} + \text{CH}_3\text{ONa}$   
 S:  $\text{C}_6\text{H}_5\text{CH} = \text{CHCl} + \text{CH}_3\text{ONa}$

- Both R and S
- Both P and R
- Both Q and S
- Both P and Q

**Ans. d**

**Sol.** Primary alkyl halides/benzyl halides reacts with alkoxide/phenoxide through  $\text{S}_{\text{N}}2$  mechanism gives ethers.

Vinyl and aryl halides least reactive towards  $\text{S}_{\text{N}}1$

18. The steps involved in the conversion of propan-2-ol to propan-1-ol are in the order


- Dehydration, addition of HBr, heating with aq. KOH
- Heating with  $\text{PCl}_5$ , heating with alc. KOH, acid catalysed addition of water
- Heating with  $\text{PCl}_5$ , heating with alc. KOH, hydroboration oxidation
- Dehydration, addition of HBr in presence of peroxide, heating with alc. KOH

**Ans. c**

19. Which of the following is the strongest base?

- $\text{CH}_3\text{COO}^-$
- $\text{Cl}^-$
- $\text{OH}^-$
- $\text{CH}_3\text{O}^-$


**Ans. d**



The product 'P' is

- 
- 
- 
- 

**Ans. c**





21. Which of the following has the lowest boiling point?  
 a)  $\text{CH}_3\text{CH}_2\text{OH}$                       b)  $\text{CH}_3 - \text{CH}_2 - \text{NH}_2$   
 c)  $\text{CH}_3 - \text{O} - \text{CH}_3$                       d)  $\text{HCOOH}$

**Ans. c**

22. The carbonyl compound that does not undergo aldol condensation is  
 a) Acetone  
 b) Di chloro acetaldehyde  
 c) Tri chloro acetaldehyde  
 d) Acetaldehyde

**Ans. c**

**Sol.** Aldehydes and ketones containing alpha hydrogens will undergo aldol condensation



**Ans. b**

24. Hinsberg's reagent is  
 a)  $(\text{CH}_3\text{CO})_2\text{O}$  / pyridine  
 b)  $\text{C}_6\text{H}_5\text{SO}_2\text{Cl}$   
 c)  $\text{C}_6\text{H}_5\text{SO}_2\text{NH}_2$   
 d)  $\text{CH}_3\text{COCl}$  / pyridine

**Ans. b**

25. Which one of the following vitamins is not stored in adipose tissue?  
 a) A                      b)  $\text{B}_6$                       c) D                      d) E

**Ans. b**

26. Hypothyroidism is caused by the deficiency of  
 a) Vitamin B-12                      b) Adrenalin  
 c) Thyroxine                      d) Glucocorticoid

**Ans. c**

27.  $\text{C}_1\text{-C}_4$  glycosidic bond is NOT found in  
 a) Maltose                      b) Sucrose  
 c) Lactose                      d) Starch

**Ans. b**

28. Which of the following polymer has strongest intermolecular forces of attraction?  
 a) Neoprene                      b) Terylene  
 c) Polythene                      d) Polystyrene

**Ans. b**

29. Which of the following monomers can undergo condensation polymerization?  
 a) Styrene                      b) Glycine  
 c) Isoprene                      d) Propene

**Ans. b**

30. A food additive that acts as an antioxidant is  
 a) BHA                      b) Saccharin  
 c) Sugar syrup                      d) Salt

**Ans. a**

31. Which of the following is not related to drug-enzyme interaction?  
 a) Allosteric site                      b) Antagonist  
 c) Co-enzymes                      d) Enzyme inhibitor

**Ans. b**

32. 0.4 g of dihydrogen is made to react with 7.4 g of dichlorine to form hydrogen chloride. The volume of hydrogen formed at 273K and 1 bar pressure is  
 a) 9.08L                      b) 4.54L                      c) 90.8L                      d) 45.4L

**Ans. b**

33. With regard to photoelectric effect, identify the correct statement among the following  
 a) Energy of  $e^-$  ejected increases with the increase in the intensity of incident light  
 b) Number of  $e^-$  ejected increases with the increase in the frequency of incident light  
 c) Number of  $e^-$  ejected increases with the increase in work function  
 d) Number of  $e^-$  ejected increases with the increase in the intensity of incident light

**Ans. d**

34. The last element of the p-block in 6<sup>th</sup> period is represented by the outer most electronic configuration

- a)  $7s^2 7p^6$   
b)  $5f^{14}6d^{10}7s^2 7p^5$   
c)  $4f^{14}5d^{10}6s^2 6p^4$   
d)  $4f^{14}5d^{10}6s^2 6p^6$

**Ans. d**

35. The conjugate base of  $NH_3$  is

- a)  $NH_4^+$     b)  $NH_4OH$     c)  $NH_2OH$     d)  $NH_2^-$

**Ans. d**

36. A gas mixture contains 25% He and 75%  $CH_4$  by volume at a given temperature and pressure. The percentage by mass of methane in the mixture is approximately\_\_\_\_\_

- a) 75%    b) 25%    c) 92%    d) 8%

**Ans. c**

37. The percentage of s-character in the hybrid orbitals of nitrogen in  $NO_2^+$ ,  $NO_3^-$  and  $NH_4^+$  respectively are

- a) 33.3%, 50%, 25%    b) 33.3%, 25%, 50%  
c) 50%, 33.3%, 25%    d) 25%, 50%, 33.3%

**Ans. c**

38. The formal charge on central oxygen atom in ozone is

- a) -1    b) 0    c) +2    d) +1

**Ans. d**

39. When the same quantity of heat is absorbed by a system at two different temperatures  $T_1$  and  $T_2$ , such that  $T_1 > T_2$ , change in entropies are  $\Delta S_1$  and  $\Delta S_2$  respectively. Then

- a)  $\Delta S_1 < \Delta S_2$     b)  $\Delta S_1 = \Delta S_2$   
c)  $S_2 > S_1$     d)  $\Delta S_2 < \Delta S_1$

**Ans. a**

**Sol.** 
$$\Delta S = \frac{q}{T}$$

q is same (constant)

$$\therefore \Delta S \propto \frac{1}{T}$$

40. The oxidation number of nitrogen atoms in  $NH_4NO_3$  are

- a) +5, +5    b) -3, +5    c) +3, -5    d) -3, -3

**Ans. b**

41. A Lewis acid 'X' reacts with  $LiAlH_4$  in ether medium to give a highly toxic gas. This gas when heated with  $NH_3$  gives a compound commonly known as inorganic benzene. The gas is

- a)  $B_2O_3$     b)  $B_2H_6$     c)  $B_3N_3H_6$     d)  $BF_3$

**Ans. b**

42. The oxide of potassium that does not exist is

- a)  $K_2O$     b)  $KO_2$     c)  $K_2O_2$     d)  $K_2O_3$

**Ans. d**

43. The metal that produces  $H_2$  with both dil HCl and NaOH (aq) is

- a) Zn    b) Mg    c) Ca    d) Fe

**Ans. a**



**Sol.** Amphoteric metals can react with both acids and bases.

44. Which of the following is NOT a pair of functional isomers?

- a)  $C_2H_5OC_2H_5$  and  $C_3H_7OCH_3$   
b)  $CH_3CH_2OH$  and  $CH_3OCH_3$   
c)  $CH_3CH_2NO_2$  and  $H_2NCH_2COOH$   
d)  $CH_3COOH$  and  $HCOOCH_3$

**Ans. a**

45. Identify 'X' in the following reaction



**Ans. b**

46. Which of the following is NOT a green house gas?

- a) CFC    b)  $CO_2$     c)  $O_2$     d)  $NO_2$

**Ans. c**

47. A metal exists as an oxide with formula  $M_{0.96}O$ . Metal M can exist as  $M^{+2}$  and  $M^{+3}$  in its oxide  $M_{0.96}O$ . The percentage of  $M^{+3}$  in the oxide is nearly  
a) 8.3%    b) 4.6%    c) 5%    d) 9.6%

**Ans. a**

**Sol.**  $M_{0.96}O$

No. of  $M^{+2}$  ions = x

No. of  $M^{+3}$  ions =  $0.96 - x$

Total positive charges = Total negative charge (in magnitude)

$$x(2) + (0.96 - x)(3) = 1(2)$$

$$2x + 2.88 - 3x = 2$$

$$-x = 2 - 2.88$$

$$\therefore x = 0.88$$

$$\begin{aligned} \text{No. of } M^{+3} \text{ ions} &= 0.96 - 0.88 \\ &= 0.08 \end{aligned}$$

$$\begin{aligned} \text{Percentage of } M^{+3} &= \frac{0.08}{0.96} \times 100 \\ &= 8.33\% \end{aligned}$$

48. A metal crystallises in face centred cubic structure with metallic radius  $\sqrt{2}A^\circ$ . The volume of the unit cell (in  $m^3$ ) is  
a)  $4 \times 10^{-10}$     b)  $6.4 \times 10^{-29}$   
c)  $4 \times 10^{-9}$     d)  $6.4 \times 10^{-30}$

**Ans. b**

**Sol.** For FCC

$$\text{Atomic radius (r)} = \frac{\sqrt{2}a}{4}$$

$$\sqrt{2} \times 10^{-10} = \frac{\sqrt{2}a}{4}$$

$$a = \frac{4 \times \sqrt{2} \times 10^{-10}}{\sqrt{2}}$$

$$a = 4 \times 10^{-10} \text{ m}$$

$$\begin{aligned} \text{Volume of unit cell} &= a^3 \\ &= (4 \times 10^{-10})^3 \\ &= 64 \times 10^{-30} \\ &= 6.4 \times 10^{-29} \text{ m}^3 \end{aligned}$$

49. Silicon doped with gallium forms  
a) n-type semiconductor  
b) both n and p type semiconductor  
c) an intrinsic semiconductor  
d) p-type semiconductor

**Ans. d**

50. The pair of electrolytes that possess same value for the constant (A) in the Debye - Huckel - Onsager equation,  $\lambda_m = \lambda_m^\circ - A\sqrt{C}$  is  
a)  $MgSO_4, NaSO_4$     b)  $NH_4Cl, NaBr$   
c)  $NaBr, MgSO_4$     d)  $NaCl, CaCl_2$

**Ans. b**

51. Which of the following pair of solutions is isotonic?  
a) 0.01M  $BaCl_2$  and 0.015M  $NaCl$   
b) 0.001M  $Al_2(SO_4)_3$  and 0.01 M  $BaCl_2$   
c) 0.001M  $CaCl_2$  and 0.001M  $Al_2(SO_4)_3$   
d) 0.01M  $BaCl_2$  and 0.001M  $CaCl_2$

**Ans. a**

**Sol.** When solute particle concentration is same then they are isotonic

52. Solute 'X' dimerises in water to the extent of 80%. 2.5g of 'X' in 100g of water increases the boiling point by  $0.3^\circ C$ . The molar mass of 'X' is [ $K_b = 0.52 K \text{ kg mol}^{-1}$ ]  
a) 13    b) 52    c) 65    d) 26

**Ans. d**

**Sol.**  $i = 1 + \alpha \left( \frac{1}{n} - 1 \right)$

$$i = 1 + 0.8 \left( \frac{1}{2} - 1 \right)$$

$$i = 1 - 0.4 = 0.6$$

$$\Delta T_b = k_b \times \frac{W}{m} \times \frac{100}{W(\text{gm})} \times i$$

$$0.3 = 0.52 \times \frac{2.5}{m} \times \frac{1000}{100} \times 0.6$$

$$\begin{aligned} \text{Molar mass of x (m)} &= \frac{0.52 \times 2.5 \times 10 \times 0.6}{0.3} \\ &= 26 \end{aligned}$$


53. Given  $E_{Fe^{+3}/Fe^{+2}}^\circ = +0.76V$  and  $E_{I_2/I^-}^\circ = +0.55V$ . The equilibrium constant for the reaction taking place in galvanic cell consisting of above two electrodes is  $\left[ \frac{2.303RT}{F} = 0.06 \right]$   
a)  $1 \times 10^7$     b)  $1 \times 10^9$     c)  $3 \times 10^8$     d)  $5 \times 10^{12}$

**Ans. a**

**Sol.**  $E_{Fe^{+3}/Fe^{+2}}^\circ = +0.76$  (cathode)

$$E_{I_2/I^-}^\circ = +0.55 \text{ (Anode)}$$

$$\begin{aligned} E_{\text{cell}}^\circ &= E_C^\circ - E_A^\circ \\ &= 0.76 - 0.55 = 0.21 \end{aligned}$$



$$E_{\text{cell}}^0 = \frac{0.059}{n} \log k_c$$

$$0.21 = \frac{0.059}{2} \log k_c$$

$$\log k_c = 7$$

$$k_c = 10^7$$

54. If an aqueous solution of NaF is electrolyzed between inert electrodes, the product obtained at anode is

- a) F<sub>2</sub>      b) H<sub>2</sub>      c) Na      d) O<sub>2</sub>

**Ans. d**

55. In which of the following cases a chemical reaction is possible ?

- a) ZnSO<sub>4(aq)</sub> is placed in a copper vessel  
b) AgNO<sub>3</sub> solution is stirred with a copper spoon  
c) Conc. HNO<sub>3</sub> is stored in a platinum vessel  
d) gold ornaments are washed with dil HCl

**Ans. b**

56. The time required for 60% completion of a first order reaction is 50 min. The time required for 93.6% completion of the same reaction will be

- a) 100 min      b) 83.8 min  
c) 50 min      d) 150 min

**Ans. d**

**Sol.** 60% completion

$$K = \frac{2.303}{t} \log \frac{[R_0]}{[R]}$$

$$K = \frac{2.303}{50} \log \frac{100}{40}$$

$$K = \frac{2.303}{50} \times 0.397$$

93.6% completion

$$K = \frac{2.303}{t} \log \frac{[R_0]}{[R]}$$

$$\frac{2.303}{50} \times 0.397 = \frac{2.303}{t} \log \frac{100}{6.4}$$

$$t = 150 \text{ min}$$

57. For an elementary reaction 2A+3B→ 4C+D the rate of appearance of C at time 't' is 2.8x10<sup>-3</sup> mol L<sup>-1</sup>S<sup>-1</sup>. Rate of disappearance of B at 't' will be

a)  $\frac{4}{3}(2.8 \times 10^{-3}) \text{ mol L}^{-1} \text{ S}^{-1}$

b)  $\frac{3}{4}(2.8 \times 10^{-3}) \text{ mol L}^{-1} \text{ S}^{-1}$

c)  $2(2.8 \times 10^{-3}) \text{ mol L}^{-1} \text{ S}^{-1}$

d)  $\frac{1}{4}(2.8 \times 10^{-3}) \text{ mol L}^{-1} \text{ S}^{-1}$

**Ans. b**

**Sol.**  $-\frac{1}{3} \frac{d(B)}{dt} = +\frac{1}{4} \frac{d(C)}{dt}$   
 $-\frac{d(B)}{dt} = +\frac{3}{4} \frac{d(C)}{dt}$   
 $= \frac{+3}{4} (2.8 \times 10^{-3}) \text{ mol L}^{-1} \text{ S}^{-1}$

58. The rate constant of a reaction is given by  $k = P Z e^{-E_a/RT}$  under standard notation. In order to speed up the reaction, which of the following factors has to be decreased ?

- a) Z      b) Both Z and T  
c) E<sub>a</sub>      d) T

**Ans. c**

59. A sol of AgI is prepared by mixing equal volumes of 0.1M AgNO<sub>3</sub> and 0.2M KI, which of the following statement is correct ?

- a) Sol obtained is a negative sol with NO<sub>3</sub><sup>-</sup> adsorbed on AgI  
b) Sol obtained is a positive sol with Ag<sup>+</sup> adsorbed on AgI  
c) Sol obtained is a positive sol with K<sup>+</sup> adsorbed on AgI  
d) Sol obtained is a negative sol with I<sup>-</sup> adsorbed on AgI

**Ans. d**

60. During Adsorption of a gas on a solid

- a) ΔG<0, ΔH<0, ΔS<0  
b) ΔG>0, ΔH>0, ΔS>0  
c) ΔG<0, ΔH<0, ΔS>0  
d) ΔG<0, ΔH>0, ΔS>0

**Ans. a**